

ЧЕСКИХ /

40%

80 %

[4, 5].

50

ГОЛОВ-

(CPS — Categorical
CPS

Pain Scale).

: 1)

; 2)

; 3)

; 4)

; 5)

ЛОБНО-ВИСОЧНОЙ
n. trigeminus

n. glossopharyngeus n. vagus).

» 17

«
CPS,

» — 19 14

— 9,
— 2;
— 6,

— 6,
: 5;
— 8,
: —

4, — 8,

— 2.

1. Ю.К., Ю.В., В.Н.
нейроиммунологические

// — 2000.

7, 1. — С. 97-100.

2. Г.Н.

— М.: , 1997. — 352 .

3. А.П.

— М.: , 1999. — 448 .

4. Шифман . . . , HELLP-

— М.: , 2002. — 430 .

5. Cuningham F.G., Twickler D. Cerebral edema complicating eclampsia 7/ Am. J. Obstet. Gynecol. — 2000. — V. 182, 1. — P. 94-100. □

PECULATES OF HEADACHE IN PREGNANT WITH PREECLAMPSY

A. Shevchenko; National Medical Academy of Post-Graduate Education named after P. Shypick, Ukraine

Summary. In the article there are results of evaluations of headache particularity in 50 pregnant persons with various severity of preeclampsy with using of Pain Scale. Increase of a threshold of irritation to pain with sensitive adaptation in pregnant with preeclampsy of light and middle severity is established as well as decrease of a threshold of irritation to pain with sensitive unadaptation to pain in the case of severe preeclampsy. It must be taken in attention during carrying out of a clinical and diagnostic estimation of severity of preeclampsy.

Key words: headache, pregnancy, preeclampsy.